



The asymptotic solutions for boundary value problem to a convective diffusion equation with chemical reaction near a cylinder

Abstract

The work deals with a boundary value problem for a quasilinear partial elliptical equation. The equation describes a stationary process of convective diffusion near a cylinder and takes into account the value of a chemical reaction for large Peclet numbers and for large constant of chemical reaction. The quantity the rate constant of the chemical reaction and Peclet number is assumed to have a constant value. The leading term of the asymptotics of the solution is constructed in the boundary layer as the solution for the quasilinear ordinary differential equation. In this paper, we construct asymptotic expansion of solutions for a quasilinear partial elliptical equation in the boundary layer near the cylinder.

Keywords

convective diffusion equation, the method of matched asymptotic expansions, the diffusion boundary layer, the saddle point, the stream function, quasilinear parabolic degenerate equation, the stability condition for difference scheme.

N. V. Maksimova^a and R. G. Akhmetov^b

N. V. Maksimova^a and R. G. Akhmetov^b

^a M.Sc: Bashkortostan State Pedagogical University, Oktyabr'skoy Revolutsii str.3a, Ufa, 450000, Russia.

^b Professor: Bashkortostan State Pedagogical University, Oktyabr'skoy Revolutsii str.3a, Ufa, 450000, Russia.

1 INTRODUCTION

The stationary convective diffusion equation in the presence of a bulk chemical reaction is given by (e.g., see [1, 2])

$$\Delta U = Pe(\bar{V}, \nabla) \cdot U + k_v F(U), \quad (1.1)$$

$$U = 1 \text{ at } r = 1; U \rightarrow 0 \text{ when } r \rightarrow \infty, \quad (1.2)$$

where

$$\bar{V} = (V_\gamma, V_\theta, 0), \quad V_\gamma = \frac{1}{r} \frac{\partial \psi}{\partial \theta}, \quad V_\theta = -\frac{\partial \psi}{\partial r}, \quad (1.3)$$

$$\psi(r, \theta) = \left(r - \frac{1}{r} \right) \sin \theta \quad (1.4)$$

is the stream function [3], r and θ are polar coordinates, Δ is the Laplace operator, Pe is the Peclet number, and k_ν is parameter depending on the chemical reaction rate. The angle θ is measured relative to the free-stream direction.

Problems analogous to (1.1) and (1.2), and a broader class of problems, were considered in [1,2], [4-6, 9, 10, 21, 22, 24]. In the absence of chemical reaction, problem (1.1) and (1.2) was analyzed in [4, 5] by the method of matched asymptotic expansions [7, 8, 23]. It is well known (see, for example, [2, Chapter 5, (6.1)-(6.3)]) that, in the limit cases $Pe \gg 1$, $k_\nu = const$; $Pe = const$ and $k_\nu \gg Pe$, the solution to problem (1.1) and (1.2) is simplified.

In the case when the volume chemical reaction of the first order ($F(u) = u$) the asymptotics of solution in all space outside the drop was constructed in [9]. In study, the number $\mu = k_\nu/Pe$ is assumed to have a constant value.

It is assumed that $F(C)$ is continuous and

$$F: R^1 \rightarrow R^1, F(0) = 0, F'(0)=0, 0 < F''(C), \quad (1.5)$$

and the asymptotic is

$$F(u) = u^2 + F_3 u^3 + F_4 u^4 + F_5 u^5 + O(u^6) \text{ for } u \rightarrow 0. \quad (1.6)$$

2 THE DIFFUSION BOUNDARY LAYER

In this report the quantity $\mu = k_\nu/Pe$ is assumed to have a constant value. In this case, all terms in Eq. (1) are similar in order of magnitude in the neighborhoods of saddle points. The small parameter $\varepsilon = (Pe)^{-1/2}$ is introduced for convenience, and Eq. (1) is rewritten as

$$\varepsilon^2 \Delta u - \frac{1}{r} \left(\frac{\partial u}{\partial r} \frac{\partial \psi}{\partial \theta} - \frac{\partial u}{\partial \theta} \frac{\partial \psi}{\partial r} \right) - \mu F(u) = 0. \quad (2.1)$$

When $\varepsilon = 0$ the Eq. (1) has the saddle points $O_1(1, \pi)$ and $O_2(1, 0)$ and the equation is equivalent the dynamical system.

The asymptotic expansions (AE) of the solution in the diffusion boundary layer was considered in a earlier study [11]. This solution was continued up to the front stagnation point $O_1(1, \pi)$ (up to the line $\theta = \pi$). The natural variables in the diffusion boundary layer are $t = \varepsilon^{-1}(r - 1)$, θ . The AE of the solution $u(t, \theta, \varepsilon)$ is sought as

$$u(t, \theta, \varepsilon) = u_0(t, \theta) + \varepsilon u_1(t, \theta) + \dots \quad (2.2)$$

From (2.1), (2.2) and (1.1) – (1.4), in variables t, θ , determining $u_0(t, \theta)$ in the domain $0 < \theta < 2\pi, 0 < t$, we obtain the problem

$$\frac{\partial^2 u_0}{\partial r^2} - 2t \cos \theta \frac{\partial u_0}{\partial r} + 2 \sin \theta \frac{\partial u_0}{\partial \theta} - \mu F(u_0) = 0, \tag{2.3}$$

$$u_0(0, \theta) = 1; \quad u_0(t, \theta) \rightarrow 0 \text{ as } t \rightarrow \infty. \tag{2.4}$$

The asymptotics of the solution to the problem (2.3), (2.4) function $u_0(t, \theta)$ as $\theta \rightarrow \pi$ is [11]

$$u_{0,0}(t) + O((\pi - \theta)^2 \exp(-\delta t^2)),$$

where $u_{0,0}(t) = O(\exp(-\delta t^2)), \delta > 0$.

3 THE ASYMPTOTICS $u_0(t, \theta)$ AS $\theta \rightarrow 0$.

The asymptotics of the function $u_0(t, \theta)$ as $t \rightarrow 0$ is sought in the view

$$V_0(t) + O(\theta^2), \tag{3.1}$$

where the function $V_0(t)$ is constructed [12] for small μ as the solution for the problem

$$L V_0 - \mu F(V_0) = V_0''(t) - t V_0'(t) - \mu F(V_0(t)) = 0 \tag{3.2}$$

$$V_0(0) = 1, V_0(t) = O(1) \text{ as } t \rightarrow \infty. \tag{3.3}$$

Theorem 1. Let $F(u)$ satisfies conditions (1.5), (1.6) and $\mu = const$, then at $t \rightarrow \infty$ the solution of the equation (3.2) asymptotics holds

$$V_0(t) = \frac{c_{01}}{\mu \ln t + C} + \frac{c_{02}}{(\mu \ln t + C)^2} + \frac{c_{03}}{(\mu \ln t + C)^3} + \dots + \ln(\mu \ln t + C) \left\{ \frac{c_{12}}{(\mu \ln t + C)^2} + \frac{c_{13}}{(\mu \ln t + C)^3} + \dots \right\} + \ln^2(\mu \ln t + C) \left\{ \frac{c_{23}}{(\mu \ln t + C)^3} + \dots \right\} + \dots \tag{3.4}$$

where

$$c_{0,1} = 1, \quad c_{0,2} = const, \quad c_{1,2} = -c_{0,1}^3 F_3, \quad c_{2,3} = c_{1,2}^2 (3 - 2c_{0,1})^{-1},$$

$$c_{1,3} = \frac{1}{3 - 2c_{0,1}}(3F_3 c_{0,1}^2 c_{1,2} + 2c_{2,3} + 2c_{0,2} c_{1,2}), \quad c_{0,3} = \frac{1}{3 - 2c_{0,1}}(c_{1,3} + c_{0,1}^4 F_4 + c_{0,2}^2 + 3F_3 c_{0,1}^2 c_{0,2}), \dots$$

The idea of the proof is similar to works [13, 14]. Let us search the function $V_0(t)$ in the form of the sum

$$V_0(t) = V_n(t) + w(t), \tag{3.5}$$

where

$$V_n(t) = \sum_{i=0}^{n-1} \ln^i(\mu \ln(t) + C) \sum_{k=i+1}^n \frac{c_{i,k}}{(\mu \ln(t) + C)^k}.$$

Substituting sum (3.5) into equation (3.2), we obtain the problem

$$Lw - \mu(F(w + V_n) - F(V_n)) = H_{n-1}(t), \tag{3.6}$$

$$w(t) \rightarrow 0, w'(t) \rightarrow 0, \text{ for } t \rightarrow \infty, \tag{3.7}$$

where $H_n(t) = O(\ln t)^{-n-1+\delta}$, δ -sufficiently small.

Let's consider the problem

$$w'' - t w' - \mu F'(V_n)w = h(t, V_n, w), \tag{3.8}$$

$$w(t) \rightarrow 0, w'(t) \rightarrow 0, \text{ for } t \rightarrow \infty, \tag{3.9}$$

where problem (3.6), (3.7) is equivalent to a problem (3.8), (3.9) and

$$h(t, V_n, w) = g(t, V_n, w) + H_{n-1}(t), \quad g(t, V_n, w) = \mu(F(w + V_n) - F(V_n) - F'(V_n)w), \tag{3.10}$$

$$g(t, V_n, w) = O(w^2)$$

For construction the solution $w(x)$ of the problem (3.8), (3.9) we obtain integral equation

$$w(t) = - \int_t^\infty W^{-1}(s) \phi_1(t)\phi_2(s) - \phi_1(s)\phi_2(t) h(s, V_n, w) ds, \tag{3.11}$$

where $\phi_1(t), \phi_2(t)$ are linearly independent solutions to the linear homogeneous equation:

$$w'' - t w' - \mu F'(V_n)w = 0, \tag{3.12}$$

$W(t) = \exp(t^2/2)$ is the Wronskian.

We have asymptotics for $\phi_1(t), \phi_2(t)$, using the results of the works [15, 16]

$$\varphi_1(t) = (\mu \ln t + C)^{-2} [1 + O(\ln t)^{-1+\delta}] \quad \text{for } t \rightarrow \infty \tag{3.13}$$

$$\varphi_2 = e^{\frac{t^2}{2}} t^{-1} (\mu \ln t + C)^2 [1 + O(\ln t)^{-1+\delta}] \quad \text{for } t \rightarrow \infty \tag{3.14}$$

where $0 < \delta$ - is small. Such solutions $\phi_1(t), \phi_2(t)$ of the equation (3.12) exists.

For $\phi_1(t)\phi_2(s) - \phi_1(s)\phi_2(t)$ we find estimate

$$\begin{aligned} \varphi_1(t)\varphi_2(s) - \varphi_1(s)\varphi_2(t) &= e^{\frac{s^2}{2}} s^{-1} (\mu \ln s + C)^2 (\mu \ln t + C)^{-2} [1 + O(\ln t)^{-1+\delta}] + O(\ln s)^{-1+\delta} - \\ &- e^{\frac{t^2}{2}} t^{-1} (\mu \ln t + C)^2 (\mu \ln s + C)^{-2} [1 + O(\ln s)^{-1+\delta}] + O(\ln t)^{-1+\delta} . \end{aligned} \tag{3.15}$$

We proceed by applying the method of successive approximations.

$$w_{n+1}(t) = - \int_t^\infty W^{-1}(s) [\phi_1(t)\phi_2(s) - \phi_1(s)\phi_2(t)] h(s, V_n, w_n) ds \tag{3.16}$$

We choose $w_0 \equiv 0$,

$$w_1(t) = - \int_t^\infty W^{-1}(s) [\phi_1(t)\phi_2(s) - \phi_1(s)\phi_2(t)] H_{n-1}(s) ds \tag{3.17}$$

From (3.6), (3.13) – (3.17) it is find estimate

$$|w_1| \leq M(\ln t)^{-n+\delta}, \tag{3.18}$$

then by formulas (3.10), (3.13) – (3.18) we have

$$\begin{aligned}
 & \left| w_2(t) - w_1(t) \right| \leq \left| -\int_t^\infty W^{-1}(s) \varphi_1(t)\varphi_2(s) - \varphi_1(s)\varphi_2(t) g(V_n, w_1) ds \right| \leq \\
 & \leq -\int_t^\infty W^{-1}(s) \varphi_1(t)\varphi_2(s) - \varphi_1(s)\varphi_2(t) g'(V_n, \bar{w}_1) w_1 ds \leq \mu MK(\ln t)^{-2n+1+2\delta} \leq \frac{M}{2}(\ln t)^{-n+\delta}, \quad t \gg 1, \quad (3.19) \\
 & \left| g(V_n, w_2) - g(V_n, w_1) \right| \leq g'(\bar{w}, V_n) |w_2 - w_1| \leq \mu MK(\ln t)^{-2n+2\delta}
 \end{aligned}$$

From (3.19) we obtain $|w_3(t) - w_2(t)| \leq \frac{M}{2^2}(\ln t)^{-n+\delta}$, $\delta > 0$ and $\forall n \geq 3$
 $|w_{n+1}(t) - w_n(t)| < \frac{M}{2^n}(\ln t)^{-n+\delta}$.

There exist $M > 0$ that for solution of the equation (3.11) inequality is hold $|w(t)| \leq 2M(\ln t)^{-n+\delta}$.

4 NUMERICAL SOLUTION AND FINDING THE CONSTANT c .

We rewrite Eq. (3.2) in the form of the system

$$\begin{cases} v_0'(t) = z(t) \\ z'(t) = tz(t) + \mu \cdot F(v_0(t)). \end{cases} \quad (4.1)$$

Consider system (4.1) on the interval $[0, X_0]$, Following [17, 18], we first discuss the stability conditions for the explicit Euler scheme

$$\begin{cases} v_{n+1} = v_n + hz_n \\ z_{n+1} = z_n + h[x_n z_n + \mu F(v_n)]. \end{cases} \quad (4.2)$$

Replacing $F(v_n)$ by the sum $F(v_0) + F'(v_0)(v_n - v_0)$ and assuming that t_0, v_0 , and z_0 are known and $t_n = t_{n-1} + h$, we find a solution to difference scheme (4.2).

The stability condition for difference scheme (4.2) is fulfilled [17-19] if $h \cdot \mu \cdot F'(t) \ll 1, h < 0, |hX_0| < 1$.

This implies that one should take X_0 and integrate backwards (i. e., with increments $h < 0$) in the interval $[0, X_0]$. The initial conditions at the point X_0 the form

$$v_0(X_0) = V_0, \quad z(X_0) = Z_0, \quad (4.3)$$

where V_0, Z_0 are found from (3.4)

$$V_0(t) = \frac{c_{01}}{\mu \ln t + C} + \frac{c_{02}}{(\mu \ln t + C)^2} + \frac{c_{03}}{(\mu \ln t + C)^3} + \ln(\mu \ln t + C) \left(\frac{c_{12}}{(\mu \ln t + C)^2} + \frac{c_{13}}{(\mu \ln t + C)^3} \right),$$

$$Z_0 = \frac{-\mu c_{01}}{t(\mu \ln t + C)^2} - \frac{\mu c_{02}}{t(\mu \ln t + C)^3} - \frac{\mu c_{03}}{t(\mu \ln t + C)^4} + \frac{\mu}{t(\mu \ln t + C)} \left(\frac{c_{12}}{(\mu \ln t + C)^2} + \frac{c_{13}}{(\mu \ln t + C)^3} \right) - \frac{\mu \ln(\mu \ln x + C)}{t} \left(\frac{2c_{12}}{(\mu \ln t + C)^3} + \frac{3c_{13}}{(\mu \ln t + C)^4} \right).$$

For example, according to (1.6) let $F(u) = \ln^2(1 + u)$.

The results of the numerical analysis of the problem (4.1), (4.3) for $F(u) = \ln^2(1 + u)$ are (for $\mu \in [0.5, 2]$, $c_{0,2} = 1$)

$$\begin{aligned} \mu = 0.5, c = 1.7216, z(0) = -0.1173; \quad \mu = 1, c = 1.3943, z(0) = -1.3943; \\ \mu = 1.5, c = 1.0452, z(0) = -0.2994; \quad \mu = 2, c = 0.6737, z(0) = -0.3747. \end{aligned}$$

This research was supported by the Presidents Grant for Leading Scientific Schools of the Russian Federation, project no. NSh - 6249.2010.1, and by the Federal Task Program "Research and educational professional community of innovation Russia" (contract 02.740.11.0612).

5 CONCLUSION

The problem considered in this work describes the phenomenon convective diffusion in the neighborhood the cylinder taking into account chemical reaction, a streamlined cross-flow of ideal fluid. In the vicinity of the cylinder there are few boundary layers. In this work the problem was investigated in the diffusion boundary layer. The leading term the asymptotic solution in the vicinity of the saddle point corresponding to the point of the liquid dripping with a cylinder has the form

$$V_0 = \frac{c_{01}}{\mu \ln t + C} + O(\ln(\mu \ln t + C)(\mu \ln t + C)^{-2})$$

at $t \rightarrow +\infty$ where $t = (r - 1)Pe^{1/2}$.

A similar structure of the asymptotics (up to a linear change of variable t) takes place in the diffusion boundary layer of convection-diffusion problem about the drop, streamlined flow of a viscous incompressible fluid.

Were previously considered the problem of convective diffusion in the neighborhood a spherical particle and a cylinder in the flow of a viscous incompressible fluid. In these cases, there are also a few boundary layers. If we compare the results obtained in the diffusion boundary layer, the structure of the asymptotics in the vicinity of the saddle point about spherical particle is simpler than those considered in this work and have the form [17].

$$v = c_0 + \frac{c_1}{\tau} + O(\tau^{-2}), \tau \rightarrow +\infty, \tau = Pe^{1/3}(r - 1).$$

A similar structure the asymptotics takes place in the flow around a cylinder cross-flow of a viscous incompressible fluid [26].

This work can be extended taking into account thermal conductivity of the same article [22]. Mathematical problems arising in this work and require further study in the internal boundary layers, containing the singular points of saddle type, there are also problems of hydrodynamics. Similar problems have been studied in [20], [25].

References

1. Levich V.G. Physicochemical Fluid Dynamics. Fizmatgiz. Moscow.: 1959 [in Russian].
2. Gupalo Yu. P., Polyanin A.D., Ryazantsev Yu. S. Mass and Heat Transfer between Reacting Particles and the Flow // Nauka. Moscow.: 1985 [in Russian].
3. Happel J., Brenner H. Low Reynolds Number Hydrodynamics. Englewood Cliffs. Prentice-Hall. 1965.
4. Sih P.H. and Newman J. Mass Transfer to the Rear of a Sphere in Stokes Flow // Int. J. Heat Mass Transfer. 1967. V. 10. P. 1749-1756.
5. Akhmetov R.G. An Asymptotic Solution to the Problem of Convection-Diffusion around a Sphere // Comput. Math. Math. Phys. V.38. № 5. P. 771-776. [Zh. Vychisl. Mat. Mat. Fiz. 1998. V. 38. № 5. P. 801-806.].
6. Akhmetov R. G. Asymptotic Behavior of the Solution to a Convection-Diffusion Problem with Distributed Chemical Reaction // Comput. Math. Math. Phys. 2002. V. 42. № 10. P. 1600—1608. [Vychisl. Mat. Mat. Fiz. 2002. V. 42. № 10. P. 1538-1546].
7. Van Dyke M.D. Perturbation Methods in Fluid Mechanics. Academic. New York.: 1964.
8. A. M. Il'in, Matching of Asymptotic Expansions of Solutions of Boundary Value Problems. Translations of Mathematical Monographs, Providence, RI, Amer. Math. Soc. 1992, V. 102.
9. Zhivotyagin A. F., Influence of a Homogeneous Chemical Reaction on the Distribution of Concentration in a Diffusion Wake of a Droplet // Vestn. Mosk. Gos. Univ. Ser. 1. Mat. Mekh. 1980. № 6. P. 73-78.
10. Chapman S.J., Lawry J.M.H., Ockendon J.R. Ray theory for high-Peclet-number convection-diffusion // SIAM J. Appl. Math. , 1999. Vol. 60, No.1, pp. 121-135.
11. Akhmetov R.G. Asymptotics of Solution for a Problem of Convective Diffusion with Volume Reaction Near a Spherical Drop // Proceedings of the Steklov Institute of Mathematics, Suppl. 1, 2003, pp. S8-S12.
12. Akhmetov R.G. The asymptotic expansions of the solution for the boundary value problem to a convective diffusion equation with volume chemical reaction near a spherical drop // Commun Nonlinear Sci Numer Simulat 16 (2011), pp. 2308-2312.
13. Fedoruk M V, WKB method for nonlinear second order equation, Journ vychisl. math. and math. phys. 26, (1986), 196-210/ (in Russian)
14. Kalyakin L A, Justification of Asymptotic Expansion for the Principal Resonance Equation, Proc. of the Steklov Inst. of Math. Suppl.1, (2003), S108-S122.
15. Olver F., Asymptotics and special functions. Academic press. New York. 1974.
16. Fedoruk M V, Asymptotic methods for ordinary differential equation. Nauka, Moscow, 1983. (in Russian)
17. Akhmetov R. G. Asymptotic Behavior of the Solution to the Convective Diffusion Problem in the Wake of a Particle // Comput. Math. Math. Phys. 2006. V. 46. № 5. P. 796—809. [Published in Zhurnal Vychislitel. Mat. Mat. Fiz. 2006. V. 46. № 5. P. 834-847].
18. Rakitskii Yu.V., Ustinov S.M., and Chernorutskii I.G., Numerical Methods for Solving Stiff Systems (Nauka, Moscow, 1979)[in Russian].
19. Babenko K. I., Fundamentals of Numerical Analysis (NITs, Regularnaya I khaoticheskaya dinamika, Moscow/Izhevsk, 2002) [in Russian].

20. Babenko, K.I., Vvedenskaya, N.D., and Orlova, M.G., Calculation of the Stationary Flow of a Viscous Fluid around a Circular Cylinder, *Zh. vych. mat. mat. _z.*, 1975, vol. 15, no.1, pp. 183-196.
21. *Polyanin and V. F. Zaitsev Handbook of Nonlinear Partial Differential Equations*, Chapman & Hall/CRC Press, Boca Raton–London, 2004.
22. Polyanin A.D. Exact solutions of nonlinear sets of equations of the theory of heat and mass transfer in reactive media and mathematical biology // *Theoretical Foundations of Chemical Engineering* 2004. V 38. N 6. P. 622-635.
23. J. Awrejcewicz, V.A. Krysko, *Introduction to Asymptotic Methods*. Chapman and Hall, CRC Press, Boca Raton, New York 2006.
24. Frolovskaya O., Nir A., Lavrenteva O.M., “Stationary regimes of axisymmetric thermal wake interaction of two buoyant drops at low Reynolds and high Peclet number”, *Physics of Fluids*, 18:7 (2006), 072103.
25. Baidulov V.G., Matyushin P.V., Chashechkin Yu.D. Evolution of the diffusion-induced flow over a sphere submerged in a continuously stratified fluid // *Fluid Dyn.* 2007. V. 42(2). P. 255-267.
26. Akhmetov R. G. Asymptotics of solutions of convection-diffusion problem about a cylinder with a chemical reaction // XVIII International conference "Mathematics. Economics. Education. "VI International Symposium "Fourier series and their applications." An interdisciplinary seminar on "Information and communication technology." Proceedings. Rostov n / D. Univ SFU. 2011. P. 79 – 84.

